

EMBERIZA QUADRICOLOR.



CHARACTER GENERICUS.

Rostrum conicum.

Mandibulae basi deorsum a se invicem distantes et inferiore lateribus inflexo-coarctata, superiore angustiore.

Lin. Syl. Nat. p. 308.

CHARACTER SPECIFICUS, &c.

EMBERIZA viridis, capite colloque cæruleis, cauda cum tectricibus abdomineque superiore rubris, pectore abdomineque infimo suscep- centibus.

Lath. Ind. orn. p. 417.

EMBERIZA QUADRICOLOR.

Lin. Syl. Nat. Gmel. p. 886.

Formosissimam aviculam depinximus quæ in insula Java præcipue conspicitur, et Americanæ speciei *Ciris* diætae admodum affinis est. In tabula magnitudine vera exprimitur.



THE
VARIEGATED EMBERIZA.



GENERIC CHARACTER.

Bill conical : Mandibles separating a little from the base downwards : the sides of the lower mandible bending rather inwards.

SPECIFIC CHARACTER, &c.

Green EMBERIZA with blue head and neck ; the tail, coverts, and upper part of the abdomen red ; the breast and lower part of the abdomen brownish.

Red-Rumped Bunting.

Lath. syn. 2. p. 208.

Le QUADRICOLOR.

Buff. ois. 3. p. 467.

Gros-bec de Java.

Pl. enl. 101. f. 2.

This beautiful bird is principally found in the island of Java, and is extremely nearly allied to the American species called *Emberiza Ciris*. The plate represents it in its natural size.